

TITLE: TENSOR ANALYZING POWERS IN THE  $^1\text{H}(\bar{d}, pp)n$  REACTION AT 16 MeV  
I. THE SYMMETRIC, CONSTANT-RELATIVE-ENERGY CONFIGURATIONS

AUTHOR(S): F. D. Correll, G. G. Ohlsen, Ronald E. Brown, R. A. Hardekopf,  
Nelson Jarmie, P. Schwandt, and P. Doleschall

SUBMITTED TO: Fifth International Symposium on Polarization  
Phenomena in Nuclear Physics, Santa Fe, NM,  
August 11-15, 1980

MASTER

## DISCLAIMER

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, product, or process disclosed or represented herein. Reference herein to any specific commercial product, process, or service does not necessarily constitute or imply its endorsement, recommendation, or approval by the United States Government or any agency thereof. Views and opinions expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

University of California



**LOS ALAMOS SCIENTIFIC LABORATORY**

Post Office Box 1663 Los Alamos, New Mexico 87545

An Affirmative Action/Equal Opportunity Employer

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos Scientific Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Tensor Analyzing Powers in the  $^1\text{H}(\vec{d},pp)n$  Reaction at 16 MeV  
 I. THE SYMMETRIC, CONSTANT-RELATIVE-ENERGY CONFIGURATIONS\*

F. D. Correll, G. G. Ohlsen, Ronald E. Brown, R. A. Hardekopf  
 and Nelson Jarmie  
 Los Alamos Scientific Laboratory, Los Alamos, NM 87545 USA

P. Schwandt  
 Indiana University, Bloomington, IN 47401 USA

P. Doleschall  
 Central Research Institute for Physics, H-1525 Budapest, Hungary

Most kinematically complete three-nucleon breakup experiments are designed to explore certain limited regions of phase space where the physical observables are expected to be especially sensitive to the detailed nature of the reaction dynamics. One interesting group of kinematic configurations, which first attracted attention as a result of Faddeev calculations performed by Kloeet and Tjon<sup>1</sup> and which were first studied experimentally by van Oers<sup>2</sup>, are characterized by equal polar angles of the two identical, detected particles and equal relative energies between all pairs of particles. In such configurations, a single breakup amplitude is expected to be dominant,<sup>3</sup> and this amplitude is believed to be particularly sensitive to the short-range characteristics of the nucleon-nucleon (NN) interaction.

We have measured the tensor analyzing powers  $A_{xx}$  and  $A_{yz}$  in the  $^1\text{H}(\vec{d},pp)n$  reaction at 16 MeV for a series of these "symmetric, constant-relative-energy" (SCRE) configurations. The formalism used in deriving analyzing powers from data on reactions with three particles in the final state has been described in Ref. 4, and we have adopted the symmetric choice for the y axis as described in that reference. Here, we report our experimental results and compare them with the results of recent Faddeev calculations using a separable NN potential.

Laboratory kinematic parameters for the configurations studied are listed in Table I, where quantities related to the detected protons are labelled by the subscripts 1 and 2. For each of the configurations shown, the polar angles of the protons are equal, and their azimuthal angles, which are measured from the x axis, are symmetric with respect to the y axis. The relative energy of each pair of nucleons is 1.55 MeV.

TABLE I. Laboratory kinematics of Configurations studied. Particles 1,2: protons; particle 3: neutron

$\alpha$ (deg)	$E_1 = E_2$ (MeV)	$\theta_1 = \theta_2$ (deg)	$\phi_1$ (deg)	$\phi_2$ (deg)	$E_3$ (MeV)
0	6.51	20.2	0.0	180.0	0.75
25	6.34	21.2	13.7	166.3	1.11
45	5.95	23.0	22.2	157.8	1.88
90	4.59	28.4	30.0	150.0	4.60
110	3.94	30.4	28.5	151.5	5.91
135	3.24	32.0	22.2	157.8	7.31
150	2.93	32.4	16.1	163.9	7.92
165	2.74	32.6	8.5	171.5	8.31
180	2.67	32.7	0.0	180.0	8.44

\*Work supported by the U. S. Department of Energy.

All of these configurations share the property that they can be generated by a rotation of the equilateral triangle formed in the c.m. system by the relative-momentum vectors of the final-state nucleon pairs. The rotation axis is fixed in the plane of the triangle and is perpendicular to the beam direction. That configuration in which the plane of the triangle contains the beam axis and in which the proton momenta are forward directed is taken to define a rotation angle  $\alpha$  of  $0^\circ$ . For this configuration, and for the one with  $\alpha=180^\circ$ , the corresponding laboratory geometries are coplanar; for other rotation angles, noncoplanar geometries result.

The experiments were performed by bombarding a  $1.2\text{-mg/cm}^2$ , rotating  $\text{CH}_2$  target with 16-MeV polarized deuterons. The target was located at the center of a large, cubical scattering chamber and was viewed by two  $\Delta E$ - $E$  solid-state detector telescopes fitted with circular collimator apertures that subtended angles of  $1.5^\circ$  at the target. Each telescope was mounted so that it could be rotated independently about two perpendicular axes to accommodate both coplanar and noncoplanar geometries. Beam intensities on target of 20-150 nA were used, and beam polarizations, measured using the quench-ratio technique, were typically in the range 0.75-0.80.

A slightly modified version of the three-spin-state method<sup>5</sup> was used to derive the analyzing powers from reaction yields measured separately with the incident deuteron beam in each of three m-states (+1, 0, -1). To minimize the effect of target deterioration upon the measured analyzing powers, the m-state sequence for the yield measurements was alternated between (+1, 0, -1) and (-1, 0, +1). The deuteron spin axis always lay in the horizontal plane and its polar angle  $\beta$  with respect to the beam direction was set to either  $45^\circ$  or  $90^\circ$ , depending upon the observable to be measured. The scattering chamber was also rotated about the beam axis into different orientations for measuring different observables.<sup>4</sup> For the two coplanar geometries,  $A_{xx}$  was measured with  $\beta=90^\circ$ ; for each of the noncoplanar geometries, the analyzing power combinations  $A_{yz}-\frac{1}{2}A_{xx}$  and  $-A_{yz}-\frac{1}{2}A_{xx}$  were measured with  $\beta=45^\circ$ , and the individual analyzing powers  $A_{xx}$  and  $A_{yz}$  were extracted later.

Standard fast NIM electronics and a multiple-input CAMAC ADC were used for the data acquisition. For each event analyzed,  $\Delta E$  and  $E$  signals from both telescopes and a TAC signal derived from the  $\Delta E$  detectors were digitized and stored in the MODCOMP IV acquisition computer. Coincident breakup protons were mass-identified on line and their two-dimensional energy-sharing spectrum  $E_1$  vs  $E_2$  was constructed. The locus of kinematically allowed breakup events was clearly evident for each geometry studied. There were usually few background counts in the vicinity of the equal-proton-energy point on each locus that corresponded to the c.m. configuration of interest (the "SCRE point").

The experimental data were analyzed in increments around the kinematic loop, using a computer program that constructed circular summing bins whose centers were equally spaced around the loop and assigned to each bin the counts in those channels of the  $E_1$  vs  $E_2$  array that fell within its circumference and were closer to its center than to the centers of its neighbors. This procedure was applied

to the individual m-state yields and the analyzing powers were calculated from the projected data.

The analyzing powers  $A_{xx}$  and  $A_{yz}$  in the region of the SCRE point on each locus studied were evaluated by averaging the data for the three nearest summing bins, and the results are plotted in Fig. 1 vs the c.m. rotation angle  $\alpha$ . The arc length over which the data were averaged ranged from 1.8 MeV for the smallest loci ( $\alpha=110^\circ$  and  $135^\circ$ ) to 4.2 MeV for the largest ( $\alpha=0^\circ$ ). However, the variation of the analyzing powers around the loci was generally slow and smooth, so that this procedure is believed to have reduced the statistical uncertainties in the quoted analyzing powers without significantly affecting their numerical values. Also shown in Fig. 1 are the predictions for these quantities from some recent Faddeev calculations made with a rather complete separable NN interaction.<sup>6</sup> In general, the agreement between the measured and calculated values seems fairly good, although there is some evidence that the measured values of

$A_{yz}$  tend to be slightly smaller than the calculations predict, whereas for  $A_{xx}$  the opposite appears to be true.




Fig. 1. Tensor Analyzing Powers  $A_{yz}$ ,  $A_{xx}$  vs cm Rotation Angle,  $\alpha$ . Data: Data; Curves: Faddeev calculations with S,P,D and  $^3S_1$ - $^3D_1$ .

1. W. M. Kloet and J. A. Tjon, Nucl. Phys. **A210**, 380 (1973).
2. W. T. H. van Oers, in Proc. 7th Int. Conf. on Few Body Problems in Nuclear and Particle Physics, ed. A. N. Mitra et al. (North-Holland, Amsterdam, 1976) p. 746. See also: D. I. Bonbright et al., Phys. Rev. **C20**, 879 (1979).
3. J. Bruinsma and R. van Wageningen, Nucl. Phys. **A282**, 1 (1977); J. H. Stuivenberg and R. van Wageningen, Nucl. Phys. **A304**, 141 (1978).
4. G. G. Ohlsen, R. E. Brown, F. D. Correll, and R. A. Hardekopf, Nucl. Instrum. Methods, to be published. Also contribution to this symposium.
5. G. G. Ohlsen and P. W. Keaton, Jr., Nucl. Instrum. Methods, **109**, 41 (1973).
6. F. D. Correll, Ronald E. Brown, G. G. Ohlsen, R. A. Hardekopf, Nelson Jarmie, and P. Doleschall, contribution to this symposium.